| Chapter 15 HW #1: Due 12/4/19 Complete both free response questions. One will be graded. Show al work. Box and clearly label all final answers $CH_3NH_2(aq) + H_2O(l) \rightleftharpoons CH_3NH_3^*(aq) + OH^*(aq) K_b = 4.4 \times 10^{-4}$ Methylamine, CH_3NH_3, is a weak base that reacts with water according to the equation above. A student obtains as 50.0 mL sample of a methylamine solution and determines the pH of the solution to be 11.77. (a) Write the expression for the equilibrium constant, K_b , for methylamine of the methylamine solution before it reacted with water and equilibrium was established. The 50.0 mL sample of the methylamine solution is titrated with an HCl solution of unknown concentration. The equivalence point of the titration is reached after a volume of 36.0 mL of the HCl solution is added. The pH of the solution at the equivalence point askes place during the titration. (c) Calculate the concentration of the HCl solution used to titrate the methylamine. (b) Calculate the initial molar expression at the equivalence point of the titration is reached after a volume of 36.0 mL of the HCl solution is added. The pH of the solution that represents the reaction that takes place during the titration. (c) Calculate the concentration of the HCl solution used to titrate the methylamine. (b) Calculate the concentration of the HCl solution used to titrate the methylamine. (c) Calculate the concentration of the HCl solution of the titration of the threat of the titration of the threat of the titration of the titrat | Name AP Chemis | stry | | | | | | | | | | |
|---|---|----------------------|--------|----------|-------|---------------------|-------|---------|---------|-------|----------------|----|
| Methylamine, CH ₃ NH ₂ , is a weak base that reacts with water according to the equation above. A student obtains a 50.0 mL sample of a methylamine solution and determines the pH of the solution to be 11.77. (a) Write the expression for the equilibrium constant, <i>K_b</i> , for methylamine. (b) Calculate the molar concentration of OH in the 50.0 mL sample of the methylamine solution. (c) Calculate the initial molar concentration of CH ₃ NH ₂ (<i>aq</i>) in the solution before it reacted with water and equilibrium was established. (a) Write the reacted with water and equilibrium was established. (b) Calculate the initial molar concentration of CH ₃ NH ₂ (<i>aq</i>) in the solution before it reacted with water and equilibrium was established. (a) Write the net-ionic equation that represents the reaction that akes place during the titration. (b) Calculate the net-ionic equation that represents the reaction that akes place during the titration. (c) Calculate the concentration of the HCl solution used to titrate the methylamine. (d) Write the expression for the equilibrium constant, <i>K_b</i> , for methylamine. (e) Calculate the concentration of the HCl solution used to titrate the methylamine. (f) Using the axes provided, sketch the titration curve that results | | espon | se qı | uestio | ons. | One | will | be g | rade | d. Sl | now | a |
| sample of the methylamine solution. (c) Calculate the initial molar concentration of CH ₃ NH ₂ (aq) in the solution before it reacted with water and equilibrium was established. The 50.0 mL sample of the methylamine solution is titrated with an HCl solution of unknown concentration. The equivalence point of the titration is reached after a volume of 36.0 mL of the HCl solution is added. The pH of the solution at the equivalence point is 5.98. (d) Write the net-ionic equation that represents the reaction that takes place during the titration. (e) Calculate the concentration of the HCl solution used to titrate the methylamine. (f) Using the axes provided, sketch the titration curve that results | Methylamine, CH_3NH_2 , is a weak base that reacts with water accord a 50.0 mL sample of a methylamine solution and determines the pH (a) Write the expression for the equilibrium constant, K_b , for methylamine. | ling to the of the s | ne equ | ation | above | | udent | obtai | ns | | | |
| The 50.0 mL sample of the methylamine solution is titrated with an HCl solution of unknown concentration. The equivalence point of the titration is reached after a volume of 36.0 mL of the HCl solution is added. The pH of the solution at the equivalence point is 5.98. (d) Write the net-ionic equation that represents the reaction that takes place during the titration. (e) Calculate the concentration of the HCl solution used to titrate the methylamine. (f) Using the axes provided, sketch the titration curve that results | sample of the methylamine solution. (c) Calculate the initial molar concentration of CH ₃ NH ₂ (aq) in the solution before it reacted with water and equilibrium was | + | | | | | | | | | | + |
| solution is added. The pH of the solution at the equivalence point s 5.98. (d) Write the net-ionic equation that represents the reaction that akes place during the titration. (e) Calculate the concentration of the HCl solution used to titrate the methylamine. (f) Using the axes provided, sketch the titration curve that results | The 50.0 mL sample of the methylamine solution is titrated with an HCl solution of unknown concentration. The equivalence point | рН | | | | | | | | | | |
| (e) Calculate the concentration of the HCl solution used to titrate the methylamine. (f) Using the axes provided, sketch the titration curve that results (g) Calculate the concentration of the HCl solution used to titrate to the methylamine. (g) 5 10 15 20 25 30 35 40 45 45 45 45 45 45 45 45 45 45 45 45 45 | solution is added. The pH of the solution at the equivalence point s 5.98. | . | | | | | | | | | | |
| 1) Using the axes provided, sketch the thration curve that results | (e) Calculate the concentration of the HCl solution used to titrate he methylamine. | 0 | | 5 1 | | 5 2 | 20 2 | 25 3 | 30 3 | 5 4 | 0 4 | 45 |
| | | mivalen | ce noi | int of t | | | | Cl Add | ed (mL) |) | | |
| | | uivalen | ce poi | int of t | | | | Cl Add | ed (mL) |) | | |
| | | uivalen | ce poi | int of t | | | | Cl Addo | ed (mL) | | | |
| | | uivalen | ce poi | int of t | | | | Cl Addo | ed (mL) | | | |
| | | uivalen | ce poi | int of t | | | | Cl Add | ed (mL) | | | |

$C_6H_5NH_2(aq) + H_2O(l) \leftarrow \rightarrow C_6H_5NH_3^+(aq) + OH^-(aq)$

Aniline, a weak base, reacts with water according to the reaction represented above.

- (a) Write the equilibrium constant expression, K_b , for the reaction represented above.
- (b) A sample of aniline is dissolved in water to produce 25.0 mL of a 0.10 M solution. The pH of the solution is 8.82. Calculate the equilibrium constant, K_b , for this reaction.
- (c) The solution prepared in part (b) is titrated with 0.10 *M* HCl. Calculate the pH of the solution when 5.0 mL of the acid has been added.
- (d) Calculate the pH at the equivalence point of the titration in part (c).
- (e) The pK_a values for several indicators are given below. Which of the indicators listed is most suitable for this titration? Justify your answer.

| Indicator | p <i>Ka</i> |
|-----------------------|-------------|
| Erythrosine | 3 |
| Litmus | 7 |
| Thymolphthalein | 10 |
| 1 ilyinoipiitiiaiciii | 10 |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |