

Chapter 5 Collected AP Exam Free Response Questions 1980 - 2010**1982 - #5**

- (a) From the standpoint of the kinetic-molecular theory, discuss briefly the properties of gas molecules that cause deviations from ideal behavior.
- (b) At 25°C and 1 atmosphere pressure, which of the following gases shows the greatest deviation from ideal behavior? Give two reasons for your choice.
- CH₄ SO₂ O₂ H₂
- (c) Real gases approach ideality at low pressure, high temperature, or both. Explain these observations

1984- #4b

- Give a scientific explanation for the following observations. Use equations or diagrams if they are relevant.
- (b) Burning coal containing a significant amount of sulfur leads to "acid rain."


1990 - #2

- A mixture of H₂(g), O₂(g), and 2 milliliters of H₂O(l) is present in 0.500-liter rigid container at 25°C. The number of moles of H₂ and the number of moles of O₂ are equal. The total pressure is 1,146 millimeters of mercury. (The equilibrium vapor pressure of pure water is 24 millimeters mercury.)
- The mixture is sparked, and H₂ and O₂ react until one reactant is completely consumed.
- (a) Identify the reactant remaining and calculate the number of moles of the reactant remaining.
- (b) Calculate the total pressure in the container at the conclusion of the reaction if the final temperature is 90°C. (The equilibrium vapor pressure of water at 90°C is 526 millimeters mercury.)
- (c) Calculate the number of moles of water present as vapor in the container at 90°C.

1993 - #9

- Observations about real gases can be explained at the molecular level according to the kinetic molecular theory of gases and ideas about intermolecular forces. Explain how each of the following observations can be interpreted according to these concepts, including how the observation supports the correctness of these theories.
- (a) When a gas-filled balloon is cooled, it shrinks in volume; this occurs no matter what gas is originally placed in the balloon.
- (b) When the balloon described in part (a) is cooled further, the volume does not become zero; rather, the gas becomes a liquid or solid.
- (c) When NH₃ gas is introduced at one end of a long tube while HCl gas is introduced simultaneously at the other end, a ring of white ammonium chloride is observed to form in the tube after a few minutes. This ring is closer to the HCl end of the tube than the NH₃ end.
- (d) A flag waves in the wind.

1994 - #3




GAS SAMPLE DATA	
Volume of sample	90.0 mL
Temperature	25° C
Atmospheric Pressure	745 mm Hg
Equilibrium Vapor Pressure of H ₂ O (25° C)	23.8 mm Hg

A student collected a sample of hydrogen gas by the displacement of water as shown by the diagram above. The relevant data are given in the following table.

- Calculate the number of moles of hydrogen gas collected.
- Calculate the number of molecules of water vapor in the sample of gas.
- Calculate the ratio of the average speed of the hydrogen molecules to the average speed of the water vapor molecules in the sample.
- Which of the two gases, H₂ or H₂O, deviates more from ideal behavior? Explain your answer.

1996 - #5



Represented above are five identical balloons, each filled to the same volume at 25°C and 1.0 atmosphere pressure with the pure gas indicated.


- Which balloon contains the greatest mass of gas? Explain.
- Compare the average kinetic energies of the gas molecules in the balloons. Explain.
- Which balloon contains the gas that would be expected to deviate most from the behavior of an ideal gas? Explain.
- Twelve hours after being filled, all the balloons have decreased in size. Predict which balloon will be the smallest. Explain your reasoning.

2002B - #2

A rigid 8.20 L flask contains a mixture of 2.50 moles of H₂, 0.500 mole of O₂, and sufficient Ar so that the partial pressure of Ar in the flask is 2.00 atm. The temperature is 127°C.

- Calculate the total pressure in the flask.
- Calculate the mole fraction of H₂ in the flask.
- Calculate the density (in g L⁻¹) of the mixture in the flask.

The mixture in the flask is ignited by a spark, and the reaction represented below occurs until one of the reactants is entirely consumed.



- Give the mole fraction of all species present in the flask at the end of the reaction.

2003 - #2

A rigid 5.00 L cylinder contains 24.5 g of $\text{N}_2(g)$ and 28.0 g of $\text{O}_2(g)$.

- (a) Calculate the total pressure, in atm, of the gas mixture in the cylinder at 298 K.
 (b) The temperature of the gas mixture in the cylinder is decreased to 280 K. Calculate each of the following.
 (i) The mole fraction of $\text{N}_2(g)$ in the cylinder
 (ii) The partial pressure, in atm, of $\text{N}_2(g)$ in the cylinder
 (c) If the cylinder develops a pinhole-sized leak and some of the gaseous mixture escapes, would the ratio

$$\frac{\text{moles of } \text{N}_2(g)}{\text{moles of } \text{O}_2(g)}$$

in the cylinder increase, decrease, or remain the same? Justify your answer.


A different rigid 5.00 L cylinder contains 0.176 mol of $\text{NO}(g)$ at 298 K. A 0.176 mol sample of $\text{O}_2(g)$ is added to the cylinder, where a reaction occurs to produce $\text{NO}_2(g)$.

- (d) Write the balanced equation for the reaction.
 (e) Calculate the total pressure, in atm, in the cylinder at 298 K after the reaction is complete.


2004 - #8d & e

Answer the following questions about carbon monoxide, $\text{CO}(g)$, and carbon dioxide, $\text{CO}_2(g)$. Assume that both gases exhibit ideal behavior.

- (d) A 1.0 mole sample of $\text{CO}(g)$ is heated at constant pressure. On the graph below, sketch the expected plot of volume versus temperature as the gas is heated.



- (e) Samples of $\text{CO}(g)$ and $\text{CO}_2(g)$ are placed in 1 L containers at the conditions indicated in the diagram below.




- (i) Indicate whether the average kinetic energy of the $\text{CO}_2(g)$ molecules is greater than, equal to, or less than the average kinetic energy of the $\text{CO}(g)$ molecules. Justify your answer.
 (ii) Indicate whether the root-mean-square speed of the $\text{CO}_2(g)$ molecules is greater than, equal to, or less than the root-mean-square speed of the $\text{CO}(g)$ molecules. Justify your answer.
 (iii) Indicate whether the number of $\text{CO}_2(g)$ molecules is greater than, equal to, or less than the number of $\text{CO}(g)$ molecules. Justify your answer.

2004B - #2


Answer the following questions related to hydrocarbons.

- (a) Determine the empirical formula of a hydrocarbon that contains 85.7 percent carbon by mass.
 (b) The density of the hydrocarbon in part (a) is 2.0 g L^{-1} at 50°C and 0.948 atm .
 (i) Calculate the molar mass of the hydrocarbon.
 (ii) Determine the molecular formula of the hydrocarbon.
 (c) Two flasks are connected by a stopcock as shown below.




The 5.0 L flask contains CH_4 at a pressure of 3.0 atm, and the 1.0 L flask contains C_2H_6 at a pressure of 0.55 atm. Calculate the total pressure of the system after the stopcock is opened. Assume that the temperature remains constant.

- (d) Octane, $\text{C}_8\text{H}_{18}(l)$, has a density of 0.703 g mL^{-1} at 20°C . A 255 mL sample of $\text{C}_8\text{H}_{18}(l)$ measured at 20°C reacts completely with excess oxygen as represented by the equation below.



Calculate the total number of moles of gaseous products formed.

2005B - #6

Consider two containers of volume 1.0 L at 298 K, as shown above. One container holds 0.10 mol $\text{N}_2(g)$ and the other holds 0.10 mol $\text{H}_2(g)$. The average kinetic energy of the $\text{N}_2(g)$ molecules is $6.2 \times 10^{-21} \text{ J}$. Assume that the $\text{N}_2(g)$ and the $\text{H}_2(g)$ exhibit ideal behavior.

- (a) Is the pressure in the container holding the $\text{H}_2(g)$ less than, greater than, or equal to the pressure in the container holding the $\text{N}_2(g)$? Justify your answer.
 (b) What is the average kinetic energy of the $\text{H}_2(g)$ molecules?
 (c) The molecules of which gas, N_2 or H_2 , have the greater average speed? Justify your answer.
 (d) What change could be made that would decrease the average kinetic energy of the $\text{N}_2(g)$ molecules in the container?
 (e) If the volume of the container holding the $\text{H}_2(g)$ was decreased to 0.50 L at 298 K, what would be the change in each of the following variables? In each case, justify your answer.
 (i) The pressure within the container
 (ii) The average speed of the $\text{H}_2(g)$ molecules

2006 - #3b


A different compound, which has the empirical formula CH_2Br , has a vapor density of 6.00 g L^{-1} at 375 K and 0.983 atm . Using these data, determine the following.

- The molar mass of the compound
- The molecular formula of the compound

2006B - #8b

Use chemical and physical principles to account for each of the following.

The inside of a metal container was cleaned with steam and immediately sealed. Later, the container imploded.

2007B - #3a

In a hydrogen-oxygen fuel cell, energy is produced by the overall reaction represented above.

- (a) When the fuel cell operates at 25°C and 1.00 atm for 78.0 minutes, 0.0746 mol of $\text{O}_2(\text{g})$ is consumed. Calculate the volume of $\text{H}_2(\text{g})$ consumed during the same time period. Express your answer in liters measured at 25°C and 1.00 atm .

2008B - #1a

Answer the following questions regarding the decomposition of arsenic pentafluoride, $\text{AsF}_5(\text{g})$.

- (a) A 55.8 g sample of $\text{AsF}_5(\text{g})$ is introduced into an evacuated 10.5 L container at 105°C .
- What is the initial molar concentration of $\text{AsF}_5(\text{g})$ in the container?
 - What is the initial pressure, in atmospheres, of the $\text{AsF}_5(\text{g})$ in the container?


2009 - #2

A student was assigned the task of determining the molar mass of an unknown gas. The student measured the mass of a sealed 843 mL rigid flask that contained dry air. The student then flushed the flask with the unknown gas, resealed it, and measured the mass again. Both the air and the unknown gas were at 23.0°C and $750. \text{ torr}$.

The data for the experiment are shown in the table below.

Volume of sealed flask	843 mL
Mass of sealed flask and dry air	157.70 g
Mass of sealed flask and unknown gas	158.08 g

- Calculate the mass, in grams, of the dry air that was in the sealed flask. (The density of dry air is 1.18 g L^{-1} at 23.0°C and $750. \text{ torr}$.)
- Calculate the mass, in grams, of the sealed flask itself (i.e., if it had no air in it).
- Calculate the mass, in grams, of the unknown gas that was added to the sealed flask.
- Using the information above, calculate the value of the molar mass of the unknown gas. After the experiment was completed, the instructor informed the student that the unknown gas was carbon dioxide (44.0 g mol^{-1}).
- Calculate the percent error in the value of the molar mass calculated in part (d).
- For each of the following two possible occurrences, indicate whether it by itself could have been responsible for the error in the student's experimental result. You need not include any calculations with your answer. For each of the possible occurrences, justify your answer.
 - Occurrence 1: The flask was incompletely flushed with $\text{CO}_2(\text{g})$, resulting in some dry air remaining in the flask.
 - Occurrence 2: The temperature of the air was 23.0°C , but the temperature of the $\text{CO}_2(\text{g})$ was lower than the reported 23.0°C .
- Describe the steps of a laboratory method that the student could use to verify that the volume of the rigid flask is 843 mL at 23.0°C . You need not include any calculations with your answer.

2009B – #3

The mass of an aqueous solution of H_2O_2 is 6.951 g. The H_2O_2 in the solution decomposes completely according to the reaction represented above. The $\text{O}_2(g)$ produced is collected in an inverted graduated tube over water at 23.4°C and has a volume of 182.4 mL when the water levels inside and outside of the tube are the same. The atmospheric pressure in the lab is 762.6 torr, and the equilibrium vapor pressure of water at 23.4°C is 21.6 torr.

- Calculate the partial pressure, in torr, of $\text{O}_2(g)$ in the gas-collection tube.
- Calculate the number of moles of $\text{O}_2(g)$ produced in the reaction.
- Calculate the mass, in grams, of H_2O_2 that decomposed.
- Calculate the percent of H_2O_2 , by mass, in the original 6.951 g aqueous sample.
- Write the oxidation number of the oxygen atoms in H_2O_2 and the oxidation number of the oxygen atoms in O_2 in the appropriate cells in the table below.

Substance	Oxidation Number of Oxygen Atoms
H_2O_2	
O_2	

- Write the balanced oxidation half-reaction for the reaction.